



SPRFMO

South Pacific Regional Fisheries Management Organisation

Jack Mackerel Working Group

Benchmark meeting report 2026

07 June 2026

22:39 PDT

Summary

1. This report summarizes the 2026 Jack Mackerel Working Group benchmark meeting. It records the main workshop decisions, agreed recommendations, unresolved technical issues, and follow-up tasks needed to support the June 2026 management strategy evaluation workshop and subsequent Scientific Committee review.

2. The benchmark meeting focused on assessment inputs, abundance-index evaluation, biological data and assumptions, assessment-model simplification, operating-model conditioning, and priorities for the MSE work programme. The report was developed from meeting notes, technical presentations, comments, and working-group discussion. Artificial intelligence tools were used extensively to help convert notes and comments into report text, with content reviewed and edited by the report authors and meeting participants.

Meeting Details

- Meeting: JMWG benchmark meeting 2026
- Date: 18-22 May 2026
- Location: Lima, Peru
- Organisation: South Pacific Regional Fisheries Management Organisation
- Working group: Jack Mackerel Working Group
- Host: Peru / IMARPE
- Chair: Jim Ianelli
- Meeting format: Primarily in person, with selected remote participation as arranged

Opening of the Meeting

3. The Chair opened the meeting and welcomed participants to the 2026 benchmark workshop. The Chair noted that the workshop has a practical purpose: to convert a large set of technical contributions into clear benchmark priorities, model-input decisions, and follow-up tasks for the JMWG and MSE process.
4. The Chair emphasized that the benchmark material contains many sensitivities, including stock structure, index selection, catchability and availability, effort creep, selectivity, recruitment assumptions, biological inputs, and operating-model conditioning. The workshop should focus on which sensitivities materially affect advice, operating-model conditioning, and MSE readiness.
5. Members were encouraged, to the extent practical, to supplement technical presentations with concise information on data extent, including years covered, spatial coverage, fleet or survey coverage, observation unit, sample sizes, treatment of missing data, and whether uncertainty is available for assessment use.
6. The Chair also noted that work sessions and report-writing time are part of the technical work of the meeting. Each session should aim to leave behind short text on decisions, unresolved issues, data requests, and recommended follow-up.

Adoption of Agenda

7. The meeting reviewed the draft agenda and agreed to organize the week around acoustic indices, CPUE indices, biological inputs, assessment-model development, projections, operating-model specifications, and final report adoption. A shortened agenda is provided in Appendix A.

Workshop Objectives

8. The main objectives of the benchmark meeting are to:
 1. Review and document candidate abundance indices and their suitability for assessment use, operating-model conditioning, and possible management-procedure input.
 2. Review biological inputs, including maturity, natural mortality, weight-at-age, length-weight relationships, and length-frequency information.
 3. Review the SC13 assessment base model and candidate simplifications for benchmark and MSE use.
 4. Identify a defensible set of assessment-model alternatives, sensitivity runs, and diagnostics.
 5. Define operating-model conditioning priorities and uncertainty dimensions for the June 2026 MSE workshop.

6. Document decisions, unresolved technical issues, and follow-up responsibilities in the meeting report.

Organization of Work

9. The meeting work should be organized around technical sessions and smaller drafting or analysis groups. Where possible, each major topic should produce:

- a short decision statement;
- a list of accepted inputs or candidate inputs;
- a list of sensitivities or diagnostics to run;
- data or documentation requests;
- text that can be inserted directly into the final workshop report.

10. For contributors who are not working directly in GitHub, use Google Docs as the collaborative drafting layer and keep this Quarto report as the source of record. The recommended workflow is described in the [Report Collaboration Workflow](#).

Key Discussion Summary

General data recommendations

11. The meeting noted that many working papers and presentations analysed important abundance-index data, but did not consistently document the extent of the data used in each analysis. This limited the group's ability to evaluate representativeness, sample support, survey coverage, and whether uncertainty was adequately propagated. Members were requested to provide concise metadata for each candidate index or supporting analysis, including years covered, spatial extent, survey or fleet coverage, observation unit, number of observations or transects, number of biological samples or tows, expansion methods, whether lengths are fork length or total length, and the uncertainty available for assessment use.

12. The catch and fishery data quality-check presentation emphasized that the JMWG has historically worked under a compressed assessment-data schedule. Members noted that the current data-submission templates and associated processing routines are pragmatic tools for meeting that schedule, and that development of Secretariat database infrastructure would not by itself resolve the immediate benchmark needs. The meeting requested that the quality-check work be posted to the `jjmData` GitHub repository, and that the group clarify how members use terms such as sampling, coverage, and sample size. In particular, the meeting requested checks on the age-length-key sample-size reporting to identify discrepancies between reported numbers of ages and numbers of length measurements.

Acoustic Indices

13. The Day 1 acoustic session combined formal SCW16 working papers with presentation-only material. The presentations covered the history of the north-central Chile acoustic survey programme, preliminary 2026 survey results, oceanographic conditions during the 2026 survey, spatio-temporal standardization of acoustic backscatter, fishery-dependent acoustic information, Peruvian acoustic assessment information, sonar applications relevant to jack mackerel and anchoveta fisheries, and acoustic information collected by fishing vessels in Peru.

14. For reporting purposes, the presentation material should be separated into four categories: formal papers tied to numbered SCW16 documents, presentation-only background, new 2026 evidence, and context or supporting methods. This distinction is important because not all Day 1 material is intended to become a direct assessment input. Some presentations provide candidate abundance indices, while others provide survey context, environmental interpretation, diagnostic information, or methodological background.

15. Presentation summaries are provided in Appendix D. Those tables should be updated as presenters confirm titles, authors, and whether each presentation should be cited as a working paper, supporting presentation, or meeting-record item.

16. For the north-central Chile acoustic survey programme, the meeting requested a summary table giving, by year where possible, the number of transects, area covered, number of trawls or biological samples used to interpret backscatter, available length-frequency information, and how biological samples were expanded. This table is needed to compare the 2024-2026 surveys against earlier years and to distinguish changes in abundance from changes in survey extent or data support. Members also requested clarification of the table presented for the 2026 survey, particularly the distinction between total survey area, area containing jack mackerel, NASC density, and biomass-related quantities.

17. The historical review of hydroacoustic surveys in north-central Chile emphasized large interannual changes in jack mackerel distribution. Participants discussed whether the recent decline and distribution in 2026 resembled earlier periods, including 1997 and the years when jack mackerel were not detected in one or more surveys. The meeting noted that the 2003-2004 period involved movement away from the survey area and that 2010 marked a return of fish to the survey area. The 2026 information suggested a different recent pattern, with limited movement from the open ocean towards the coast and an increased presence of juveniles in northern coastal areas relative to recent years.

18. The preliminary 2026 Chile acoustic results raised questions about vertical distribution, detectability near the surface, and school structure. The meeting noted that a high proportion of individuals was observed in the upper 30 m in 2025 and 2026, but that participants did not identify a survey dead-zone issue for these observations. Participants discussed whether fish close to the surface or more dispersed schools could affect detectability. The presentation indicated that the distribution of fish in 2026 compared to 2025 was more patchy, which may

render it largely undetectable by acoustics, and that acoustic-trawl operations were most effective around dawn, when jack mackerel tended to concentrate.

19. The working group had a long discussion about whether to include the most recent 2026 acoustic survey data point in the assessment runs, noting that the survey biomass estimate had declined from more than 3 million t in 2025 to a preliminary value just over 0.64 million t in 2026. Members also noted that the Chilean fleet's year-to-date catch was about 30% of the comparable catch level from the previous year. The Chair requested that members provide their best estimate of the most likely 2026 catch so that at least some indication of the apparent stock trend could be relayed before the MSE meeting and Scientific Committee review. Part of the difficulty was that the benchmark had two related but distinct goals: to improve future assessment modelling, and to generate appropriate scenarios for MSE testing. If the benchmark were being used only to improve the assessment, there would be less immediate need to resolve the 2026 treatment because the issue could be taken up at the Scientific Committee meeting later in the year. However, the need to define plausible MSE scenarios made the question more time-sensitive. One member was reluctant to conduct that model run, citing too much uncertainty in estimating 2026 catch levels, even as a sensitivity. The group noted the awkwardness of proposing an MSE under conditions that appear to be exceptional, given the apparent dispersion of jack mackerel and pending El Niño conditions.

20. The oceanographic presentation proposed that large jack mackerel may have moved offshore and dispersed in response to reduced coastal food concentration associated with ENSO-related inhibition of coastal upwelling. The meeting noted that this hypothesis should be compared directly with the acoustic observations, because some survey evidence appeared to suggest coastal or near-coastal concentrations rather than offshore displacement. The report should therefore treat the oceanographic explanation as a working hypothesis that requires reconciliation with the acoustic distribution, chlorophyll-a information, and fishery-dependent observations.

21. Two spatio-temporal acoustic standardization approaches for northern Chile were reviewed. The *sdmTMB* acoustic-density analysis based on NASC used zero-included spatio-temporal modelling and was presented as a complementary index. The meeting discussed whether the Tweedie model adequately captured the very high proportion of zeros, whether the survey design and spatial mesh could absorb abundance signal, and how uncertainty in extrapolated areas should be carried through to the final index. Participants cautioned that summing median predictions across areas may understate uncertainty if spatially extrapolated cells are weakly informed. The second northern Chile analysis also used *sdmTMB*, with acoustic information converted to biomass using biological data, and participants discussed the roles of occurrence, positive density, anisotropy, and environmental covariates. The meeting noted that these two acoustic products use related data and should be compared for complementarity, redundancy, uncertainty, and sensitivity to survey coverage. The meeting also noted that there would be an advantage to developing a synoptic acoustic survey design that covers the northern to southern parts of Chile, so that future survey information can better distinguish changes in abundance from shifts in distribution and availability.

22. On Day 3, the meeting revisited the Chile Acoustic North survey treatment. Participants noted that the spatio-temporal modelling work by INPESCA is a substantial contribution and should be acknowledged in the report. The group noted that spatio-temporal modelling of acoustic data is generally less common than spatio-temporal standardization of CPUE. The latter time series is largely affected by fishers' choice, which often disguises changes in population densities. The Chile Acoustic North survey, however, follows a design-based approach with pre-determined transects that are largely comparable across years and hence do not suffer from fisheries behaviour in the same way. It is assumed that the survey observes all relevant fish densities in the area and hence does not require interpolation or extrapolation to other areas such as those used for the CPUE standardization. The Day 3 decision was to retain the raw or design-based acoustic biomass treatment for assessment use at this stage, while using the spatio-temporal acoustic modelling as research and diagnostic information rather than as a replacement base-model index.

23. The fishery-dependent acoustic presentation for south-central Chile was viewed as potentially useful but not yet ready for direct assessment use without further work. Participants noted the high temporal resolution and lower cost of commercial-vessel acoustic information, but also discussed preferential sampling, changes in fishing behaviour, and seasonal shifts. The meeting discussed whether fishery-dependent and fishery-independent acoustic data could eventually be combined, but noted that the fishery-dependent product may be more appropriate as a diagnostic or future sensitivity until the effects of seasonality, vessel coverage, and fleet behaviour are better understood.

24. For the Chile South-Central acoustic survey time series, participants noted that significant changes in acoustic survey sampling designs have been necessary to track shifts in Chilean jack mackerel distribution. During the early part of the series, from 1997-2003, survey cruises maintained a relatively stable transect design focused on the coastal zone, approximately the first 100 nmi from Valparaiso (33 degrees S) to Corral (40 degrees S). This was followed by an extended period from 2004-2017 characterized by annual changes to transect design, driven primarily by offshore movement of jack mackerel into oceanic waters. This spatial shift is supported by commercial fleet catch distributions, which show vessels operating up to 800 nmi from the coast in search of fishing grounds. In some years the survey could not operate, while in other years operational and administrative issues may have resulted in false negatives in the acoustic data, particularly during 2009-2015.

25. These methodological changes raised concerns about comparability of survey results over time, including changes in the number of transects, longitudinal extent, latitudinal extent, and overall study area. Consequently, only a restricted period of the time series, 2001-2009, has historically been used for jack mackerel stock assessments, consistent with the SCW14 benchmark agreement. Participants noted that modifying acoustic survey design is essential for highly mobile pelagic stocks whose spatial distributions shift with changing oceanographic conditions. Fixed historical grids can bias abundance estimates when fish move beyond the original sampling frame, while adaptive sampling protocols that track density boundaries can reduce these errors. However, fluctuating survey areas also challenge the assumption of constant

catchability, making raw design-based indices difficult to use directly in population models without additional standardization. Spatio-temporal standardization approaches, including methods such as VAST, were identified as tools that could account for the varying survey footprint and generate a more robust continuous biomass index for modern assessments. Chile noted that a project is underway to improve jack mackerel hydroacoustic surveys through optimized survey design, estimation of mean acoustic density and associated uncertainty, and standardization of historical survey time series using Bayesian and/or classical approaches. The project also aims to propose standardized abundance and biomass indices derived from jack mackerel acoustic surveys, with results expected to be presented at a future SPRFMO Scientific Committee meeting for consideration in the jack mackerel stock assessment model.

26. The Peruvian acoustic discussion emphasized that the acoustic survey programme was primarily designed for anchovy, and that jack mackerel may occur outside the surveyed area in some years. Participants discussed the large decline after the 1997/1998 El Nino event, possible ecological regime changes, changes in spatial distribution from nearshore to more offshore areas, and whether the existing Peru acoustic series in the assessment adequately reflects those changes. The meeting requested Peruvian acoustic survey backscatter data and associated length composition data, including survey timing, spatial extent, transect coverage, biological sampling, length type, and uncertainty. For the new acoustic surveys in the Peruvian zone, the length-composition effective sample size was scaled to the number of survey tows that included jack mackerel. For scaling, large numbers of tows were capped at 30, and the survey vector was then rescaled to an average effective sample size of 15, consistent with other surveys of this type. The meeting also noted that summer and winter surveys may need to be treated separately because seasonal differences and contrasting within-year values may contain useful information. Day 3 follow-up items included checking that the average survey month and seasonal assignment were implemented correctly in the model runs, reviewing fits to zero observations, and obtaining survey length frequencies as soon as possible.

27. The sonar and fishing-vessel acoustic presentations provided methodological context rather than immediate assessment inputs. Sonar may improve information on presence, school behaviour, swimming direction, and fish outside the echosounder beam, but was not considered ready to provide quantitative biomass estimates for jack mackerel. Fishing-vessel acoustic data in Peru may provide useful context, but additional work is needed to understand juvenile avoidance, vessel behaviour, calibration, and how fishery-dependent acoustic observations relate to fishery-independent survey information.

28. The meeting identified a need for a dedicated analysis that compares acoustic survey extent, backscatter, biological sampling, length-frequency data, and numbers-at-age across years and regions. This analysis should include northern Chile, central-south Chile, Peru, and fishery-dependent acoustic products where available. The objective is to determine which acoustic information is suitable for direct assessment likelihood use, which is suitable for operating-model conditioning or management-procedure diagnostics, and which should remain supporting context.

CPUE Indices

29. The Day 2 CPUE session reviewed catch and fishery data quality checks, Chilean effort-creep correction, spatio-temporal Chilean CPUE standardization using `sdmTMB` and INLA-based approaches, Peruvian CPUE standardization, and offshore CPUE treatment. The discussion focused on whether candidate CPUE products are ready for assessment use, what additional diagnostics are needed, and how to document any change from the current assessment CPUE treatment.

30. During the evaluation of the modelling procedure for the Chilean South-Central (SC) fleet abundance index, benchmark participants deliberated on the most robust method to account for effort creep. Initial positions diverged, with Chile proposing to apply an informed correction factor from 2005 onwards, whereas the EU advocated for maintaining a fixed 1% compounded factor across the entire time series. Arguments suggesting continuous improvement in efficiency focused on development in gear technology, improved online communication between skippers via WhatsApp, improved echosounders and sonars, and newly available AI-derived predictions of resource density via providers such as Greenfish. Industry representatives from several member countries corroborated that these innovations increased fishing efficiency. After extensive discussion, it was agreed to proceed with Chile's proposal and apply a 1% compounded increase creep factor up to 2004, followed by a transition to the p10 correction factor from 2005 onward, where a fixed multiplicative factor of 0.669 is applied to the abundance index, as proposed by Zenteno and Paya (2026). The EU objected and the Chair noted that it could be raised again at SC but should not have a large effect on making progress with the MSE.

31. Participants also discussed whether applying the creep factor as an effort correction is equivalent to imposing a catchability change, and whether the survey/interview information used to estimate creep can be maintained regularly. The meeting noted that the artisanal fleet now accounts for a larger share of Chilean catch and may need to be considered in future effort-creep work. As a diagnostic, the meeting requested a model run with no creep correction but with a catchability step change in 2005, to evaluate whether the assessment estimates a large catchability shift without imposing the creep correction directly.

32. Chile agreed to continue surveying industry on innovations that may trigger effort-creep corrections in the future.

33. For the Chilean `sdmTMB` CPUE analysis, the meeting noted that the index was updated through 2026 and is based on set-level data. Discussion focused on haul capacity treatment, the use of a spline smoother versus vessel categories, the large imbalance in early-year data coverage, and whether true zero catches are available and should be incorporated. Participants noted that the early part of the set-level data may represent a very small fraction of total catch, raising questions about representativeness before about 2001. The meeting requested that the data table include the number and proportion of sets by year, and that the authors provide a clear explanation for why the new set-based CPUE series gives lower CPUE in earlier years with high catches than the current trip-based assessment series.

34. The INLA-based Chilean CPUE analysis provided a complementary spatio-temporal standardization. Participants discussed mesh resolution, the treatment of northern fishing activity, whether assessment fleets should be defined primarily by area or vessel behaviour, and whether CPUE weighting should be based on catch rates rather than catch. The meeting noted that the model appeared sensitive to data updates, with inclusion of 2026 data changing specifically the 2021 estimate, and that this sensitivity needs to be explained before replacing the existing assessment index, for example through an analytical retrospective. Participants also discussed fixed versus random year effects, AR(1) versus independent year effects, and the interpretation of the days-at-sea or DFP coefficient. Movement of the resource further offshore could lead to increases in days at sea, which hence may not suggest a decrease in stock size although it does result in a decline in CPUE. As such, including days at sea in a CPUE standardization may bias the outcomes. Since the INLA approach accounts for annual spatial shifts by explicitly fitting an annual spatial random field, this concern is alleviated. Previous versions of the INLA model submitted to SC12 and SC13 were compared to the updated 2026 configuration. The introduction of annual spatial random fields changed the perception of the resource dynamics, with less profound increases and decreases over the time series.

35. Comparisons between the CPUE index model based on `sdmTMB` (Paya 2026) and INLA (Vasquez and Sepulveda 2026) were presented and discussed (Figure 1). There were no significant differences between model outputs. For benchmark purposes, the INLA model (Vasquez and Sepulveda 2026) was selected. For the update of the indices for the next stock assessment, which should include data up to June 2026, the group agreed on a common model configuration:

36. $C = \text{Year} + \text{Quarter} + \text{DAS} + s(\text{hc}, 5) + s$

37. where DAS is days at sea, hc is holding capacity, and s represents spatio-temporal random effects independent of software platform.

38. The updated INLA-based Chilean CPUE retrospective analysis was reviewed using six peels. The resulting Mohn's rho was 0.0254, and participants considered the diagnostic performance favourable for assessment use.

39. The Day 3 discussion agreed to move forward with a new Chilean CPUE treatment starting in 1998, using the agreed effort-creep treatment described above. Annual CVs should be weighted by percentage coverage and should incorporate standard errors from the INLA analysis. The previous Chilean CPUE index should be removed from the base model. A sensitivity run should replace the new Chilean CPUE series with the SC13 CPUE series to evaluate the effect of removing the old CPUE treatment on historical biomass.

40. The 1998 start year was selected because earlier set-level coverage was poor and may not be representative of the fishery, particularly in years when catches were large but the proportion of catch represented in the CPUE data was low. The meeting noted that the behaviour of the fleet and the amount of data collected changed substantially after 1997, coincident with the major 1997/1998 El Nino event. For the benchmark, the base model should therefore use the new INLA set-based Chilean CPUE series truncated to start in 1998. Future updates to this index




Figure 1: CPUE indices normalized as z-scores and estimated using `sdmTMB-INLA` (GLMM_ST; Paya 2026) and `INLA` (GLMM_ST; Vasquez and Sepulveda 2026).

should be documented carefully, because substantial changes to CPUE treatment between the benchmark and MSE implementation could affect perceptions of the MSE results.

41. The agreed uncertainty treatment for the new Chilean South-Central CPUE series is summarized in Table 1. The CV weight is calculated as the square-root of annual coverage relative to the maximum coverage in the series; equivalently, the table reports the inverse multiplier ($1/CV_wt$) used to inflate the annual CV. The assessment model uses the 1998-2025 rows and the weighted (`wtd`) CV column. The 0.06 CPUE index (`CPUE+creep`) column is the effort-creep-adjusted Chilean CPUE vector used as `Chile_CPUE` in the 0.06 model input file. It applies the agreed 1% compounded fishing-power correction through 2004 and the 0.669 multiplicative adjustment from 2005 onward.

Table 1: Chilean South-Central CPUE uncertainty and 0.06 input-index table. The 1998-2025 rows were selected for the assessment-model input; the 0.06 CPUE index (`CPUE+creep`) column is the `Chile_CPUE` vector used in the 0.06 model, and the weighted (`wtd`) CV is the CV used for the assessment model.

year	mean_cpue	cpue_creep_0_06	sd_cpue	cv	wtd_cv	coverage	inv_cv_wt
1994	85.370	NA	17.484	20%	359%	0.3	17.5499
1995	95.896	NA	12.781	13%	135%	0.9	10.1325
1996	109.032	NA	13.295	12%	85%	1.9	6.9736
1997	77.243	NA	11.940	15%	105%	2.0	6.7971

year	mean_cpue	cpue_creep_0_06	sd_cpue	cv	wtd_cv	coverage	inv_cv_wt
1998	100.264	96.35176	12.860	13%	42%	8.6	3.2778
1999	91.383	86.94815	12.188	13%	38%	11.4	2.8470
2000	127.563	120.17014	13.687	11%	28%	13.8	2.5876
2001	130.360	121.58886	15.797	12%	21%	30.2	1.7492
2002	120.465	111.24762	11.277	9%	18%	24.6	1.9381
2003	138.190	126.35233	11.606	8%	18%	19.8	2.1602
2004	163.079	147.63318	12.033	7%	11%	44.9	1.4345
2005	193.942	129.74712	14.731	8%	11%	43.6	1.4558
2006	178.593	119.47840	25.286	14%	49%	7.7	3.4641
2007	148.058	99.05053	16.480	11%	28%	14.5	2.5244
2008	169.673	113.51116	10.637	6%	10%	39.1	1.5373
2009	161.031	107.72963	9.894	6%	10%	36.8	1.5846
2010	141.411	94.60366	10.989	8%	11%	46.0	1.4173
2011	89.460	59.84884	7.215	8%	14%	32.6	1.6836
2012	88.672	59.32153	9.422	11%	12%	76.5	1.0990
2013	97.089	64.95235	10.991	11%	14%	56.4	1.2800
2014	92.191	61.67549	10.589	11%	24%	21.2	2.0877
2015	91.071	60.92678	7.820	9%	13%	40.3	1.5142
2016	103.501	69.24232	10.668	10%	18%	29.2	1.7789
2017	111.113	74.33426	13.865	12%	25%	23.3	1.9914
2018	104.030	69.59604	13.952	13%	31%	16.8	2.3452
2019	153.819	102.90499	18.896	12%	18%	41.2	1.4976
2020	143.967	96.31388	16.481	11%	15%	50.9	1.3473
2021	140.068	93.70528	19.265	14%	16%	69.7	1.1514
2022	134.764	90.15704	21.881	16%	17%	80.9	1.0687
2023	146.730	98.16246	22.071	15%	16%	83.4	1.0526
2024	123.286	82.47849	14.038	11%	11%	92.4	1.0000
2025	145.421	97.28663	13.947	10%	10%	88.8	1.0201
Mean	129.614	94.83282	13.877	11%	20%	81.0	1.0681
1998-2025							

42. The Peruvian CPUE presentation covered a standardized CPUE series for 2015-2025 that combines industrial and artisanal fleets. The meeting noted this combined fleet treatment as a strength, but also discussed the interpretation of trip locations, haul-capacity and distance-to-coast smoothers, possible differences between industrial and artisanal fleet effects, and changes in Peruvian fishing operations. Members noted that jack mackerel has been used for direct human consumption in Peru since 2002, and that the artisanal fishery has accounted for a larger share of the quota since 2019. These changes, in combination with environmental impacts on the distribution of jack mackerel, led to a marked shift of the resource over the CPUE time series. This shift is currently untreated in the CPUE standardization, as efforts to account for

it were unsuccessful. The Day 3 discussion agreed that the new, shorter Peruvian CPUE series provides better information and should be preferred for the base model. The previous Peru CPUE index should therefore be removed from the base configuration, while a sensitivity that also includes the old Peru CPUE series can be considered.

43. The offshore CPUE discussion included the treatment of a catchability break in 2021 associated with movement of the offshore fleet to the north, and discussion of how El Nino effects were represented relative to annual effects. The report should document the rationale for any offshore catchability break and whether environmental effects are interpreted as catchability effects, availability effects, or residual year effects.

44. As part of the meeting request to better understand patterns in data collection and fishing-set coverage for the Chilean CPUE series, Chile provided a summary table of annual fleet and coverage characteristics (Table 2). The table includes distance from port, trip duration, catch per set, vessel hold capacity, the proportion of catch referenced by the set-level data, and the number of fishing sets available by year. This table was reviewed during a lengthy discussion about the period over which the modeled CPUE series presented by Vasquez should be used. A critical concern was low data coverage in the early period and the counter-intuitive result that the CPUE standardization model indicated low abundance during years when the largest catches were made. The Chair also considered a diagnostic figure based on the mean catch per set divided by hold capacity times the number of sets, and suggested omitting 1994, 2006, and 2026 from that diagnostic as outlying observations. The resulting normalized catch-rate pattern (Figure 2) provides a simple nominal check on the annual information in the table rather than a replacement for the CPUE standardization. The diagnostic declines from the high early values toward lower values in the early 2000s, increases through the mid- to late 2010s, and then declines in the most recent retained years. From this analysis, as well as from other statistical analyses, such as PCA, evaluated during the meeting, starting the time series in 1998 was considered most suitable given that sampling coverage was similar across years from that time point onwards.

45. The broader pairwise diagnostic summary is shown in Figure 3.

Table 2: Annual Chilean South-Central fleet and fishing-set summary used to evaluate CPUE data coverage.

Year	Distance from port (km)	Trip duration (days)	Catch per set (t)	Hold capacity (m3)	Catch represented (%)	Fishing sets (n)
1994	161.9	2.2	177.2	847	0.3	69
1995	205.6	2.0	143.5	959	0.9	268
1996	381.9	3.2	171.2	1073	1.9	397
1997	192.6	2.7	118.5	1188	2.0	468
1998	222.4	2.6	153.5	1389	8.6	883
1999	197.4	2.6	148.7	1354	11.4	893

Year	Distance from port (km)	Trip duration (days)	Catch per set (t)	Hold capacity (m3)	Catch represented (%)	Fishing sets (n)
2000	224.9	2.6	181.1	1484	13.8	832
2001	180.6	2.4	174.8	1367	30.2	2407
2002	270.5	3.4	174.4	1243	24.6	1970
2003	413.0	3.3	164.6	1228	19.8	1534
2004	447.9	3.3	225.1	1272	44.9	2578
2005	483.2	3.5	220.5	1251	43.6	2504
2006	284.8	2.5	244.6	1299	7.7	388
2007	448.7	4.1	192.0	1357	14.5	854
2008	947.0	7.2	189.6	1346	39.1	1505
2009	845.4	7.2	165.3	1375	36.8	1557
2010	970.2	9.2	151.3	1395	46.0	897
2011	769.1	8.4	95.9	1532	32.6	732
2012	253.3	3.6	125.5	1438	76.5	1293
2013	331.7	3.9	141.9	1468	56.4	853
2014	300.2	5.7	114.5	1577	21.2	471
2015	541.7	6.2	111.6	1518	40.3	901
2016	355.6	4.3	159.9	1626	29.2	531
2017	286.5	4.0	157.8	1614	23.3	449
2018	281.9	3.8	152.7	1674	16.8	455
2019	196.2	2.6	211.2	1561	41.2	823
2020	204.5	2.4	253.8	1564	50.9	1036
2021	185.0	2.5	247.6	1564	69.7	1596
2022	122.1	2.3	242.9	1585	80.9	2001
2023	136.8	1.6	252.9	1578	83.4	2379
2024	196.9	2.8	232.7	1583	92.4	5237
2025	375.0	3.8	233.1	1608	88.8	4163
2026	221.8	3.9	238.8	1584	81.0	954

Biological Inputs

46. The Day 3 biological-input discussion reviewed the Peruvian length-weight and length-frequency material and agreed that the Peruvian length frequencies should be treated as fork length. Growth parameters used with those data should therefore also be in fork length. Members noted the SPRFMO convention that length information should be reported in fork length rather than total length. The group agreed to leave the length-composition plus group at 50 cm so the Peruvian length-frequency information and growth assumptions are handled consistently in the model.




Figure 2: Normalized annual catch-rate diagnostic calculated as mean catch per set divided by hold capacity and number of fishing sets. The plot omits 1994, 2006, and 2026 and shows annual points only.




Figure 3: Pairs plot of annual Chilean South-Central fishing-trip and set-coverage variables from `fishing_trip_summary_1994_2026.csv`. Lower panels show annual points labeled by year with loess smoothers; upper panels show Pearson correlations.

47. The meeting also requested that Peru provide sample-size information for the composition data in a form that can be translated into assessment input sample sizes. Members noted that the number of fish measured would be useful in addition to the current reporting of sampling events such as boxes or trips. The group also discussed whether the SPRFMO length-bin range should be expanded to match the Peruvian 5-90 cm length bins.

48. Jim presented model runs using the updated Peruvian length-composition data. Members requested additional checks on the resulting selectivity patterns, particularly that selectivity should not become dome-shaped after 2020. The Peruvian length-weight presentation also noted that the allometric parameter b fluctuates and tends to decline sharply during El Nino events, which should be considered when interpreting biological inputs from years affected by environmental anomalies.

Far North length-composition binning and data specification

49. The updated Far North length-composition information was provided in annual rows for 1972-2025 with centimetre bins from 5 to 90 cm. The assessment input currently uses 10-50 cm length bins. The meeting reviewed cumulative length distributions and annual length-composition shapes to evaluate whether expanding the model length-bin range was necessary for the benchmark configuration.

50. The diagnostic plots show that the retained 10-50 cm interval captures nearly all of the observed length-composition mass in most years (Figure 4; Figure 5; Figure 6). Across the annual rows, the median proportion within 10-50 cm was 0.9998, and the minimum was 0.895. When the length compositions were summed across all years, 0.9986 of the total was within the retained 10-50 cm interval. The lower annual coverage years were mainly early observations and the most recent 2024 row. On this basis, the practical benchmark decision was to retain the existing 10-50 cm assessment bins for the Far North length-composition treatment rather than expanding the JJM length-bin range to 5-90 cm at this stage.

51. The annual length-composition sample-size plot was digitized from the provided figure for 1972-2025. The digitized values were treated as relative sample-size weights. The meeting noted that rescaling the Far North length-frequency sample sizes to have a maximum value of 100 would not preserve the scale used previously in the assessment. Instead, the effective sample-size vector should maintain the same mean as in the past. Consequently, the vector was rescaled to have a mean effective sample size of 30. The scaled input vector is archived in `doc/data/farnorth_sample_sizes_scaled_for_mod0.5.csv`.

52. The Season 1 and Season 2 Far North length-composition rows used in `mod0.5.dat` were also reviewed against the retained 10-50 cm assessment bins (Figure 7; Figure 8). The ridge plots preserve gaps for missing years so the temporal coverage of each seasonal series is explicit.




Figure 4: Cumulative Far North length-composition distribution after summing across all years. The curve uses the full 5-90 cm composition data, with the x-axis truncated at 60 cm. Dashed vertical lines mark the retained 10-50 cm assessment length-bin interval.




Figure 5: Cumulative Far North length-composition distributions by year. Lines show row-wise cumulative proportions from the full 5-90 cm composition data, with the x-axis truncated at 60 cm. Dashed vertical lines mark the retained 10-50 cm assessment length-bin interval.




Figure 6: Annual Far North length-composition shapes by year. Each ridge is scaled within year to emphasize modal structure; dashed vertical lines mark the retained 10-50 cm assessment length-bin interval.




Figure 7: Season 1 Far North length-composition shapes from mod0.5.dat. Each ridge is scaled within year to emphasize modal structure; dashed vertical lines mark the retained 10-50 cm assessment length-bin interval, and blank rows indicate missing years.




Figure 8: Season 2 Far North length-composition shapes from mod0.5.dat. Each ridge is scaled within year to emphasize modal structure; dashed vertical lines mark the retained 10-50 cm assessment length-bin interval, and blank rows indicate missing years.

53. After discussions regarding the representativeness of the jack mackerel (JM) abundance estimates off Peru, given that the coverage of the Peruvian acoustic surveys, which are primarily designed to assess Peruvian anchovy, does not fully match the distribution of JM, the meeting agreed to move to the new Peruvian acoustic indices as indices of availability and remove the previous Peru acoustic index from the base configuration. The associated Peruvian length-frequency information should be carried forward with the new acoustic treatment. No new Chilean Acoustic North or Chilean Acoustic South-Central index treatment was selected for the base model at this stage.

Assessment Model Development

54. The meeting reviewed Paper 01 ([SCW16-Doc01](#)), which evaluated and updated the previous assessment model from SC13. Participants noted that the paper covered a broad set of model alternatives, including single-stock and two-stock configurations, reduced-index options, selectivity alternatives, and other candidate simplifications intended to improve benchmark and MSE tractability.

55. The group also noted that several simplification suggestions in Paper 01 should be interpreted in light of the data updates agreed during the benchmark meeting. Because several abundance-index and composition series were updated or replaced during the meeting, some simplifications that were informative under the SC13 data configuration may no longer apply directly to the updated benchmark configuration. The final candidate model set should therefore reassess those simplification options after the updated input series are implemented, rather than carrying all earlier simplification conclusions forward unchanged.

56. Participants also discussed whether 2026 information should be included in the benchmark or MSE model runs. One suggested approach was to estimate total 2026 catch from year-to-date Chilean catch information and use that estimate in a sensitivity run. The group also reviewed options for Chile Acoustic South-Central data treatment, including dropping the series, retaining it as currently used, updating CVs based on geostatistical spatial coverage, transect design, and operational reliability, or retaining the series with heavier downweighting. The practical proposal carried forward was to retain the series as currently configured unless a later sensitivity is requested.

57. The group discussed diagnostics of the 0.06 model, noting that sample sizes for the Far North fleet were high. It was agreed to rescale these sample sizes to have an average of 30 over the time series. New diagnostic plots, including one-step-ahead residuals, were created and deemed valuable for model-performance evaluation. Several iterations are still needed to improve model fit. The Chair suggested allowing until Wednesday 27 May 2026 to improve model fit and double-check data inputs and model settings. The group agreed to this approach.

Operating Models and MSE Linkages

58. The working group conferred with the MSE contractor on details of the projection and scenario specifications needed to link the benchmark assessment work to the MSE process. Discussion focused in particular on how recruitment process errors would be developed for the projection runs. The contractor clarified that an ARIMA-based treatment may be preferable, with both the autocorrelation parameter (**rho**) and residual scale (**sigma**) estimated from that process. Under that treatment, **sigma** is interpreted as the innovation standard deviation: the remaining recruitment variability after accounting for autocorrelation. The working group agreed in principle with the proposed base-model configuration runs, noting that the final base-model set used for MSE conditioning will arise from the benchmark decisions.

59. The group reviewed the operating-model grid paper presented by I. Mosquera and discussed how the benchmark model set would condition the South Pacific jack mackerel MSE. Participants noted that, although preliminary 2025 information has been presented as best available science, the conditioning model uses abundance information from the beginning of 2025 and should not be interpreted as using 2025 as a complete year. Discussion of projection selectivity and reference-point assumptions accepted at the previous Scientific Committee was tabled until supporting documentation is available. The group identified an urgent need to specify transition assumptions for selectivity and weight-at-age in projections, given the strong mandate to have an MSE ready during 2026. The group recognized that these specifications are prerequisites for the contractor to advance the MSE work. The group acknowledged that any delay in providing these specifications would directly affect the contractor's ability to deliver against the agreed timeline, and that progress on the MSE is contingent on timely decisions by the group.

60. Participants also discussed projected recruitment deviations and whether terminal-year uncertainty should be propagated directly into future recruitment. The group noted the need to separate estimation uncertainty from prediction and requested that Iago examine the distribution of values for **sigma_R**, which had not been revisited since the September model. Future recruitment variability should consider whether to exclude the extremely high 1986 and 1987 recruitment events or use a more recent recruitment period. The group also noted that exceptional circumstances, including the apparent 2026 dispersion of jack mackerel, need to be considered when framing MSE scenarios and subsequent advice. Day 5 discussions revisited the estimation of recruitment deviations, as the contractor had simulated deviates in an alternative manner, including autocorrelation correction for recruitment deviates. This resulted in substantially narrower ranges of deviates, in line with recent and long-term observations. Since autocorrelation was estimated at around 0.68, it was assumed that this parameter was reliably estimated and hence the newly developed approach seemed appropriate. The group considered recent recruitment deviations most suitable for the MSE simulations.

61. The group discussed management-procedure implementation questions, including how to rank indices for management-procedure input versus assessment and conditioning use, and whether MSE scenarios should include banking and borrowing provisions approved at the

previous Commission meeting. The current MSE setup can represent banking and borrowing across all fleets, rather than by fleet, and the specific TAC-change bounds should be checked against the Commission report.

Decisions and Agreements

62. The table below consolidates the benchmark decisions and agreements reached during the meeting and identifies the follow-up work needed to translate them into model inputs, sensitivity runs, and final report text. Items are grouped by technical topic so that implementation checks can be tracked alongside the action-item list.

Table 3: Benchmark decisions and agreements. Numbered entries in the Status column identify the corresponding model reference.

No.	Decision or agreement	Topic	Follow-up needed	Status
1	Use the new Chilean CPUE series beginning in 1998.	Chile CPUE	Implement in the base model and remove the previous Chilean CPUE index.	0.01
2	Apply the Chilean effort-creep treatment with 1% annual creep through 2004 and the 0.669 adjustment beginning in 2005.	Chile CPUE / effort creep	Confirm the exact implementation in the model input and control files.	0.06
3	Weight Chilean CPUE CVs by percentage coverage and include standard errors from the INLA analysis.	Chile CPUE uncertainty	Prepare the annual uncertainty vector for assessment input.	0.05
4	Run a sensitivity replacing the new Chilean CPUE series with the SC13 CPUE series.	Chile CPUE sensitivity	Evaluate the impact of removing the old CPUE series on historical biomass.	TBD
5	Use the new short Peruvian CPUE series as the preferred base-model treatment.	Peru CPUE	Implement the new series in the base model and consider a sensitivity that also includes the old Peru CPUE series.	0.02
6	Treat Peruvian length frequencies as fork length and use growth parameters in fork length.	Peru biology / composition	Check consistency of growth, length-frequency bins, and length-weight inputs.	0.00+

No.	Decision or agreement	Topic	Follow-up needed	Status
7	Retain the 50 cm plus group for the Peruvian length-frequency and growth treatment.	Peru biology / composition	Confirm the retained plus-group treatment in the candidate model configuration.	Complete
8	Use the new Peruvian acoustic indices and remove the previous Peru acoustic index.	Peru acoustic index	Carry forward associated length-frequency information with the new acoustic treatment.	0.05
9	Retain raw/design-based Chile Acoustic North biomass for assessment use and keep spatio-temporal acoustic modelling as research and diagnostic information.	Chile acoustic indices	Acknowledge INPESCA's modelling contribution and document why model-based CPUE indicators and design-based survey indicators are treated differently.	Research
10	Do not add a new base-model treatment for Chilean Acoustic South-Central at this stage.	Chile acoustic indices	Retain as diagnostic or supporting information unless a later sensitivity is requested.	Complete
11	Estimate variable fishery selectivity for Peru catch length frequencies.	Peru fishery selectivity	Implement or test variable selectivity for Peru catch length-frequency data.	TBD
12	Use Peru survey length frequencies to estimate survey selectivity.	Peru survey selectivity	Link Peru survey length frequencies to survey selectivity estimation in the model configuration.	TBD

Benchmark Summary and Recommendations

63. Based on the 2026 benchmark workshop of the Jack Mackerel Working Group, the following points are directed to the Scientific Committee and to the upcoming MSE workshop.

- **Noted** that the benchmark reviewed and re-evaluated multiple data streams, including acoustic surveys, Chilean and Peruvian CPUE indices, biological inputs, composition data, and candidate assessment-model configurations.
- **Noted** that several abundance-index and composition series were updated or replaced during the meeting, so model simplification and MSE conditioning should be evaluated

against the benchmark data configuration rather than against the previous SC13 input set.

- **Noted** that the new Chilean CPUE treatment begins in 1998, includes the agreed effort-creep adjustment, and uses annual uncertainty weighted by data coverage with INLA model-estimated standard errors.
- **Noted** that the 2026 Chile acoustic survey and year-to-date fishery information indicate unusual recent conditions, including apparent dispersion of jack mackerel and a large decline in the north-central Chile acoustic biomass estimate relative to 2025.
- **Noted** that acoustic and CPUE data may give different recent signals and that interpretation of availability, catchability, survey coverage, and biological sampling remains important for both assessment and MSE use.
- **Recommended** using the agreed benchmark index set in the candidate base-model configuration, including the new Chilean CPUE series, the new Peruvian CPUE series, and the new Peruvian acoustic indices, while removing the previous Peru CPUE and acoustic index treatments from the base configuration.
- **Recommended** retaining the raw or design-based Chile Acoustic North biomass treatment for assessment use, while treating spatio-temporal acoustic modelling as research and diagnostic information unless a later sensitivity is requested.
- **Recommended** that the JMWG and relevant Members evaluate the feasibility of a synoptic acoustic survey covering the northern to southern parts of Chile. The group emphasized the importance of such surveys and data to distinguish abundance trends from spatial redistribution and availability changes.
- **Recommended** retaining the existing 10-50 cm Far North length-composition bins and 50 cm plus group for the benchmark configuration, using Peruvian length-frequency information as fork length and applying fork-length growth parameters.
- **Recommended** that the MSE workshop use the benchmark-derived base-model set for operating-model conditioning and explicitly document recruitment-process assumptions, including any ARIMA-based treatment of autocorrelation and innovation variance.
- **Recommended** that the JMWG develop a set of meta-rules for identifying and responding to exceptional circumstances, including cases where recent survey, fishery, environmental, or distributional information may fall outside the conditions anticipated by the operating models or management procedures.
- **Recommended** presenting near-term model sensitivities or diagnostics, where feasible, that reflect the 2026 acoustic and catch information so the Scientific Committee and MSE process have a clear indication of the apparent recent stock trend and its uncertainty.

References

64. The meeting report relied on the SCW16 working papers and supporting presentations listed below. Presentation-only material and file crosswalks are provided in Appendix D.

- SCW16-Doc01. Developments of the base SC13 model for benchmark and MSE considerations.
- SCW16-Doc02. CPUE coordination priorities and working-paper plan.
- SCW16-Doc03. Meta-analysis of CPUE papers on jack mackerel.
- SCW16-Doc04. Conditioning of an operating-model grid for South Pacific jack mackerel.
- SCW16-Doc05. Jack mackerel biology summary.
- SCW16-Doc06. Standardized northern Chile acoustic abundance index.
- SCW16-Doc07. Implementation of an informed creep correction into the Chile jack mackerel CPUE index.
- SCW16-Doc08. Interannual variability in distribution, size structure, and biomass estimated from fishery-dependent acoustics.
- SCW16-Doc09. Spatio-temporal modelling of acoustic backscatter from north-central Chile acoustic surveys using `sdmTMB`.
- SCW16-Doc10. Chilean CPUE standardization using `sdmTMB`.
- SCW16-Doc11. Jack mackerel catch and fishery data quality-control checks.
- SCW16-Doc12. Spatio-temporal modelling of Chilean CPUE using INLA.
- SCW16-Doc13. Length-weight relationship of jack mackerel in Peru.
- SCW16-Doc14. Updated length-frequency data for Peruvian jack mackerel.
- SCW16-Doc15. Standardization of CPUE for jack mackerel in Peru.
- SCW16-Doc16. Considerations on the use of the jack mackerel acoustic index in Peruvian national jurisdictional waters.
- Harbitz, A., Ona, E., and Pennington, M. (2009). The use of an adaptive acoustic-survey design to estimate the abundance of highly skewed fish populations. *ICES Journal of Marine Science*, 66, 1349-1354. <https://doi.org/10.1093/icesjms/fsp088>.
- Stenevik, E. K., Volstad, J. H., Hoines, A., Aanes, S., Oskarsson, G. J., Jacobsen, J. A., and Tangen, O. (2015). Precision in estimates of density and biomass of Norwegian spring-spawning herring based on acoustic surveys. *Marine Biology Research*, 11, 449-461. <https://doi.org/10.1080/17451000.2014.995672>.
- Thorson, J. T. (2019). Guidance for decisions using the Vector Autoregressive Spatio-Temporal (VAST) package in stock, ecosystem, habitat and climate assessments. *Fisheries Research*, 210, 143-161. <https://doi.org/10.1016/j.fishres.2018.10.013>.
- Paya, I. (2025). Update on the Chilean jack mackerel CPUE index and acoustic biomass estimates in Chile. South Pacific Regional Fisheries Management Organisation (SPRFMO). <https://sprfmo.int/assets/Meetings/02-SC/13th-SC-2025/Jack-Mackerel/SC13-JM04-Jack-Mackerel-CPUE-index-and-acoustic-biomass-in-the-south-central-Chile-up-to-2025-.pdf>.
- Vasquez, S., and Sepulveda, A. (2025). Update on CPUE standardization for the Chilean jack mackerel fishery in central-southern Chile using spatio-temporal Bayesian models. South Pacific Regional Fisheries Management Organisation (SPRFMO). <https://www.sprfmo.int/assets/Meetings/02-SC/13th-SC-2025/Jack-Mackerel/SC13-JM02-Update-on-CPUE-standardization-for-CJM-fishery-in-central-southern-Chile.pdf>.
- SCW14. (2022). 14th Scientific Committee Workshop Report: Jack Mackerel Benchmark Workshop. South Pacific Regional Fisheries Management Organisation (SPRFMO).

https://www.sprfmo.int/assets/Meetings/SC_WS/SCW14-Jack-Mackerel/SPRFMO-SC-JM-Benchmark-Workshop-2022-Report-SCW14.pdf

Appendix A. Shortened Draft Agenda

This appendix summarizes the draft agenda from SCW16-Doc00. It should be updated if the adopted meeting agenda differs materially.

Day 1: Acoustic Session

- Opening ceremony, official photo, and meeting organization.
- Historical review of hydroacoustic survey programs in north-central Chile.
- Preliminary results of the 2026 north-central Chile hydroacoustic survey.
- Oceanographic conditions during the 2026 acoustic survey.
- Spatio-temporal modeling of acoustic backscatter from the north-central Chile survey.
- Toward a standardized northern Chile acoustic abundance index.
- Opportunistic acoustic data collection from the south-central Chilean commercial fleet.
- Review of jack mackerel acoustic assessments conducted in Peru, including SCW16-Doc16.
- Use of acoustic data collected by fishing vessels for estimating abundance/biomass in Peru.
- Echogram examination and validation.
- General discussion and technical recommendations.

Day 2: CPUE Session

- Metadata by fleet and country.
- CPUE methods by fleet and country.
- Peruvian CPUE report, including SCW16-Doc15.
- Comparative analysis and parsimony across candidate indices.
- Effort-creep review and decision for stock assessment use.
- Catchability and availability blocks by abundance index.
- CPUE weighting factors for stock assessment.
- Separate versus combined index products for HCR or management-procedure use.

Day 3: Recap, Report Drafting, and Biological Inputs

- Recap of acoustic and CPUE sessions.
- Drafting of index-session report text and follow-up tasks.
- Summary of SCW16-Doc01 and benchmark-model implications.

- Review of biological assumptions used in the assessment and operating models.
- Review of Peru length-weight and length-frequency information, including SCW16-Doc13 and SCW16-Doc14.

Day 4: Projections, Recruitment Scenarios, and Operating Models

- Assessment projections and recruitment scenarios.
- Candidate assessment-model runs and sensitivity structure.
- Operating-model specifications for MSE conditioning.
- MCMC evaluations, convergence diagnostics, run-time constraints, and feasibility.

Day 5: Final Decisions and Report Adoption

- Final benchmark decisions and recommendations.
- Confirmation of model-run priorities, operating-model specifications, and follow-up tasks.
- Report drafting, review, and adoption.

Appendix B. Participants

This participant list is based on the registration information available at the start of the benchmark meeting. Email addresses are retained outside this public report.

Name	Delegation	Role	Organisation / agency	Format
Jose Ignacio Zenteno Loredo	Chile	Advisor	IFOP	In person
Aquiles Sepulveda	Chile	Advisor	INPESCA	In person
Qi Lee	Invited Expert	Invited Expert by SPRFMO	University of Washington	In person
Niels Teon Hintzen	European Union	Expert	PFA	In person
Karolina Molla Gazi	European Union	Designated Representative / HoD	Wageningen Marine Research	In person
Jim Ianelli	JMWG Chairperson	Expert / Chair	NOAA	In person
Victor Catasti	Chile	Expert	IFOP	In person
Ignacio Paya	Chile	Expert	IFOP	In person

Name	Delegation	Role	Organisation / agency	Format
Sebastian Vazquez	Chile	Expert	INPESCA	In person
Nicolas Alegria	Chile	Expert	INPESCA	In person
Nicole Mermoud	Chile	Designated Representative / HoD	SUBPESCA	In person
Ana Alegre Norza Sior	Peru	Designated Representative / HoD	Instituto del Mar del Peru	In person
Criscely Lujan Paredes	Peru	Expert	Instituto del Mar del Peru	In person
Mirian Geronimo Aparicio	Peru	Expert	Instituto del Mar del Peru	In person
Gersson Roman Amancio	Peru	Expert	Instituto del Mar del Peru	In person
Erich Diaz Acuna	Peru	Expert	Instituto del Mar del Peru	In person
Josymar Torrejon	Peru	Expert	Instituto del Mar del Peru	In person
Daniel Isaias Grados Paredes	Peru	Expert	Instituto del Mar del Peru	In person
Carlos Valdez Mego	Peru	Expert	Instituto del Mar del Peru	In person
Marissela Pozada Herrera	Peru	Expert	Instituto del Mar del Peru	In person
Mariano Sergio Gutierrez Torero	Peru	Expert	Instituto Humboldt	In person
Salvador Peraltilla Neyra	Peru	Expert	Sociedad Nacional de Pesqueria	In person
Sandra Marisol Cahuin Villanueva	Peru	Expert	Instituto del Mar del Peru	Virtual
Iago Mosqueira	Expert	Expert	WUR	Virtual
Teobaldo Dioses	Peru	Expert	Instituto del Mar del Peru (retired)	In person
Jaime Letelier	Chile	Expert	IFOP	TBD
Esteban Molina	Chile	Expert	IFOP	TBD

Name	Delegation	Role	Organisation / agency	Format
Francisco Leiva	Chile	Expert	IFOP	TBD
Javier Legua	Chile	Expert	IFOP	TBD
Jorge Castillo	Chile	Expert	IFOP	TBD

Appendix C. Document Summary by Topic

This appendix summarizes SCW16-Doc01 through SCW16-Doc16. The organization is by topic rather than document number so that related evidence can be reviewed together.

Assessment and Model Structure

Developments of the Base SC13 Model

[SCW16-Doc01](#) documents benchmark development from the SC13 JJM assessment model, including single-stock and two-stock configurations, reduced-index sensitivities, selectivity alternatives, and model diagnostics. It is the main bridge between the current assessment model and the benchmark model set.

Key report use: summarize accepted base-model structure, candidate simplifications, retained sensitivities, and any model configurations to carry into OM conditioning.

Operating Models and MSE

Operating Models for Jack Mackerel MSE

[SCW16-Doc04](#) describes the operating-model structure for jack mackerel MSE. It should be used to organize reference-set and robustness-set uncertainties, including stock hypotheses, recruitment assumptions, selectivity, catchability, and future projection structure.

Key report use: define the operating-model grid or equivalent uncertainty table that will feed the June 2026 MSE workshop.

CPUE

CPUE Coordination Priorities and Working-Paper Plan

[SCW16-Doc02](#) provides the forward-looking CPUE coordination plan for the benchmark, including candidate topics, decision points, metadata needs, and working-paper expectations.

Key report use: structure the CPUE session and make sure each candidate CPUE product is classified for assessment, OM conditioning, MP input, diagnostic use, or hold.

Meta-Analysis of CPUE Papers on Jack Mackerel

[SCW16-Doc03](#) synthesizes CPUE papers from 2022-2025. It provides historical context on CPUE methods, index conflicts, effort creep, spatial behavior, environmental effects, and recurring recommendations.

Key report use: provide background for interpreting new CPUE papers and avoid repeating previously identified unresolved issues.

Chile CPUE Effort Creep

[SCW16-Doc07](#) focuses on implementation of an informed creep correction for the Chilean jack mackerel CPUE index. It is central to decisions about fishing-power correction and whether a revised CPUE series should be used directly or treated as a sensitivity.

Key report use: document whether effort-creep treatment is accepted, treated as a sensitivity, or deferred pending additional diagnostics.

Chile CPUE with sdmTMB/SPDE

[SCW16-Doc10](#) updates the Chilean CPUE abundance index using spatio-temporal SPDE-based models with sdmTMB. It should be compared directly with other Chilean CPUE standardization approaches.

Key report use: evaluate whether the spatio-temporal CPUE product should replace, supplement, or diagnose the current assessment CPUE series.

Chile CPUE with INLA

SCW16-Doc12 standardizes the central-southern Chile CPUE fishery using hierarchical Bayesian models with INLA. It provides a complementary spatio-temporal approach to Doc10.

Key report use: compare diagnostics, uncertainty, spatial extrapolation, and annual index behavior against the sdmTMB CPUE product.

Peru CPUE

SCW16-Doc15 presents standardization of catch-per-unit-effort for jack mackerel from 2015-2025 in Peruvian national jurisdictional waters.

Key report use: evaluate whether the Peruvian CPUE product is suitable as a far-north abundance index, an assessment sensitivity, an OM-conditioning input, or a diagnostic series.

Acoustic Indices

Northern Chile Acoustic Spatio-Temporal Model

SCW16-Doc06 develops a standardized abundance index for Chilean jack mackerel using spatio-temporal modeling of acoustic survey data in northern Chile.

Key report use: evaluate acoustic index representativeness, uncertainty, spatial coverage, and suitability for assessment and OM use.

Fishery-Dependent Acoustics

SCW16-Doc08 summarizes interannual variability in distribution, size structure, and biomass estimated from fishery-dependent acoustics.

Key report use: decide whether fishery-dependent acoustic products provide a usable abundance signal or primarily diagnostic/context information.

Central-North Acoustic Density with sdmTMB

SCW16-Doc09 estimates a jack mackerel acoustic density index for central-north acoustic surveys using spatio-temporal models with sdmTMB.

Key report use: compare with Doc06 and other acoustic products to determine whether indices are complementary, redundant, or conflicting.

Peru Acoustic Index Considerations

[SCW16-Doc16](#) summarizes considerations on the use of the jack mackerel acoustic index in Peruvian national jurisdictional waters.

Key report use: classify the Peru acoustic information as a candidate index, supporting interpretation, or diagnostic evidence, depending on data extent, uncertainty, and assessment compatibility.

Biology and Composition Inputs

Biology

[SCW16-Doc05](#) summarizes biological inputs for the assessment, including natural mortality, maturity, weight-at-age, and spawning-biomass calculations.

Key report use: document accepted biological assumptions and identify sensitivity tests that should be carried into assessment or OM conditioning.

Peru Length-Weight Relationship

[SCW16-Doc13](#) presents the length-weight relationship of jack mackerel in Peru and implications for biomass estimation.

Key report use: evaluate whether Peru-specific length-weight information affects biomass conversion, biological assumptions, or regional differences in assessment inputs.

Peru Length-Frequency Update

[SCW16-Doc14](#) summarizes updated length-frequency data for the Peruvian jack mackerel far-north stock in Peruvian jurisdictional waters.

Key report use: document the composition-data evidence available for the far-north stock and decide how it should be used in assessment and stock-structure discussions.

Data Quality and Supporting Evidence

Catch/Fishery Data Quality Checks

[SCW16-Doc11](#) provides catch and fishery data quality-control checks.

Key report use: use as a support screen for catch- and fishery-dependent indices, and document any data-quality issues that affect benchmark decisions.

Appendix D. Presentation summaries

This appendix records presentation files available for the benchmark meeting. It is intended as a crosswalk between the meeting agenda, formal SCW16 papers, presentation-only material, and how each item should be used in the meeting report.

For Day 1, the presentation files are available in docs/day1 and are organized around the acoustic-session agenda.

Presentation topic	Linked paper	Main contribution	Report use
Historical review of hydroacoustic survey programs in north-central Chile	Presentation only	Survey-program history, continuity, and context for interpreting the northern Chile acoustic time series.	Background / context
Preliminary results of the 2026 north-central Chile hydroacoustic survey	Presentation only; related to Chile acoustic index papers	New 2026 acoustic information and immediate interpretation of recent survey conditions and abundance signal.	New 2026 evidence / benchmark caveat
Oceanographic conditions during the 2026 acoustic survey	Presentation only	Environmental context for interpreting the 2026 survey and potential availability or distribution effects.	Context / supporting interpretation
Spatio-temporal modeling of acoustic backscatter using sdmTMB	SCW16-Doc09	Acoustic density index estimated with spatio-temporal methods.	Formal paper / candidate index
Standardized northern Chile acoustic abundance index	SCW16-Doc06	Spatio-temporal modelling of northern Chile acoustic-survey data.	Formal paper / candidate index
Opportunistic acoustic data collection using the south-central Chilean commercial fleet	SCW16-Doc08	Fishery-dependent acoustic information on distribution, size structure, and biomass.	Formal paper / diagnostic or candidate index

Presentation topic	Linked paper	Main contribution	Report use
Review of jack mackerel acoustic assessments conducted in Peru	SCW16-Doc16	Peruvian acoustic assessment context and considerations for index use.	Formal paper / supporting evidence or candidate index
Sonar applications in Peruvian fisheries and applicability for jack mackerel	Presentation only	Methods and operational context for sonar/acoustic information in Peru and Chile.	Context / supporting methods
Acoustic data collected by fishing vessels in Peru	Presentation only	Use of fishing-vessel acoustic data for estimating abundance or biomass of anchovy and Chilean jack mackerel in Peru.	Context / supporting methods

For Day 2, the presentation files currently available in docs/day2 are organized around the CPUE and data-quality session.

Presentation topic	Linked paper	Main contribution	Report use
Implementation of an informed creep correction into the Chile Jack Mackerel CPUE Index	SCW16-Doc07	Industrial-vessel fishing-power and effort-creep correction for the Chilean jack mackerel CPUE index.	Formal paper / candidate base or sensitivity treatment
Standardization of Chilean jack mackerel CPUE fishery in central-southern Chile using Hierarchical Bayesian Models	SCW16-Doc12	INLA-based spatio-temporal standardization of the central-southern Chile CPUE fishery.	Formal paper / candidate index
Standardization of CPUE for jack mackerel in Peru	SCW16-Doc15	Updated Peruvian CPUE standardization for 2015-2025 using industrial and artisanal fleet information.	Formal paper / candidate diagnostic or future index

Presentation topic	Linked paper	Main contribution	Report use
CPUE standardisation offshore fleet	Presentation only; related to offshore CPUE assessment treatment	Offshore fleet CPUE standardization, including treatment of a 2021 catchability break and environmental or catchability interpretation.	Presentation only / model-treatment diagnostic

For Day 3, the presentation files currently available in `docs/day3` are organized around CPUE decisions, acoustic survey follow-up, and biological inputs. The appendix links to compressed PDF exports rather than PPTX or ODP files. Two biology-summary source PPTX versions are listed because the local files differ by hash. The two Peruvian length-frequency source PDF filenames listed below have identical SHA-256 hashes and appear to be duplicate copies of the same presentation; both filenames are retained here to preserve the file-version crosswalk.

Presentation topic	Linked paper	Main contribution	Report use
Jack mackerel biology summary; second source version	SCW16-Doc05	Biological-input summary for the assessment, including natural mortality, maturity, weight-at-age, and spawning-biomass calculations.	Formal paper / biological input
2026 South-Central Chile jack mackerel fishery update	Presentation only; compressed PDF export of local PPTX	Recent south-central Chile fishery information, including 2026 fishing conditions relevant to interpreting CPUE and availability.	New 2026 evidence / CPUE caveat
CPUE discussion material	Presentation only; related to SCW16-Doc15	Supplemental CPUE discussion material, including spatial-distribution and catchability considerations for Peruvian CPUE interpretation.	CPUE diagnostic / model-treatment discussion

Presentation topic	Linked paper	Main contribution	Report use
Peruvian acoustic biomass discussion	Presentation only; related to SCW16-Doc16	Follow-up discussion of Peruvian acoustic biomass information, survey coverage, zero observations, and assessment-use questions.	Acoustic-survey diagnostic / data request
Updated Peruvian jack mackerel length-frequency data	SCW16-Doc14	Updated far-north Peruvian length-frequency information and sample-size considerations for assessment composition data. Source filenames retained in the crosswalk: 03_05_Updated updated length-frequencypdf and 03_CLUJAN_length_benchmark.pdf are identical files.	Biological input / composition data

Use the following labels when carrying presentation material into the report:

Label	Meaning
Formal paper	Presentation tied to a numbered SCW16 working paper.
Presentation only	Material presented to the meeting but not currently represented by a numbered SCW16 paper.
New 2026 evidence	Recent information that may affect benchmark interpretation even if it is not a standalone paper.
Context / supporting methods	Background or method information that helps interpret candidate indices but is not itself a direct assessment input.

The diagnostic figure appendix previously appended to this meeting report has been removed. It is replaced by the [SCW16 benchmark assessment report](#), which maintains the current model sequence and diagnostic material.